

On Minkowski sums of simplices

Geir Agnarsson *

Walter D. Morris *

Abstract

We investigate the structure of the Minkowski sum of standard simplices in \mathbb{R}^r . In particular, we investigate the one-dimensional structure, the vertices, their degrees and the edges in the Minkowski sum polytope.

2000 MSC: 52B05, 52B11, 05C07.

Keywords: polytope, Minkowski sum, hyperplane.

1 Introduction and Definitions

Let $[r] = \{1, 2, \dots, r\}$. The *standard simplex* $\Delta_{[r]}$ of dimension $r - 1$ is given by

$$\Delta_{[r]} = \{(x_1, \dots, x_r) \in \mathbb{R}^r : x_i \geq 0 \text{ for all } i, x_1 + \dots + x_r = 1\}.$$

Each subset $F \subseteq [r]$ yields a *face* Δ_F of $\Delta_{[r]}$ given by

$$\Delta_F = \{(x_1, \dots, x_r) \in \Delta_{[r]} : x_i = 0 \text{ for } i \notin F\}.$$

Clearly Δ_F is itself a simplex embedded in \mathbb{R}^r . If \mathcal{F} is a family of subsets of $[r]$, then we can form the *Minkowski sum* of simplices

$$P_{\mathcal{F}} = \sum_{F \in \mathcal{F}} \Delta_F = \left\{ \sum_{F \in \mathcal{F}} x_F : x_F \in \Delta_F \text{ for each } F \in \mathcal{F} \right\}.$$

If $|F| = 2$ for all $F \in \mathcal{F}$, then the polytope $P_{\mathcal{F}}$ is called a *graphical zonotope*. Graphical zonotopes were studied by West et. al. [4], [11], but several questions about them have gone unanswered. Minkowski sums of simplices have more recently been studied by Feichtner and Sturmfels [3], and by Postnikov [9]. These later papers focus on the case when the collection \mathcal{F} is a *building set*, i.e. \mathcal{F} contains all singletons, and has the property that, for any $F_1, F_2 \in \mathcal{F}$, $F_1 \cap F_2 \neq \emptyset$ implies that $F_1 \cup F_2 \in \mathcal{F}$. This property implies that the polytope $P_{\mathcal{F}}$ is simple. Applications of Minkowski sums of simplices appear in the paper of Morton et. al. [8]. Minkowski sums of simplices have also appeared in the work of Conca [2] and of Herzog and Hibi [6], under the name transversal polymatroids.

*Department of Mathematical Sciences, George Mason University, MS 3F2, 4400 University Drive, Fairfax, VA – 22030, USA, {geir@math.gmu.edu, wmorris@gmu.edu}

Observation 1.1 *The dimension of the polytope $P_{\mathcal{F}}$ is given by $\dim(P_{\mathcal{F}}) = n - c$ where*

$$n = \left| \bigcup_{F \in \mathcal{F}} F \right| \in [r]$$

and c is the number of connected components of $\Delta_{\mathcal{F}}$, the simplicial complex with facets $\max(\mathcal{F})$.

Proof. For each $F \in \mathcal{F}$ present Δ_F by $\Delta_F = \{(x_{F;1}, \dots, x_{F;n}) \in \Delta_{[r]} : x_{F;i} = 0 \text{ for } i \notin F\}$. We then use the equation $\sum_{i \in F} x_{F;i} = 1$ for each F to obtain $x_{F;\max(F)} = 1 - \sum_{i \in F \setminus \{\max(F)\}} x_{F;i}$ and eliminate $x_{F;\max(F)}$ in the Minkowski sum. By then counting the free variables, we have the observation. \square

From the following more graph theoretic point of view we also can consider the following: Let $\Delta_1(\mathcal{F})$ be the 1-dimensional skeleton of $P_{\mathcal{F}}$.

Observation 1.2 *The dimension of the polytope $P_{\mathcal{F}}$ is given by $\dim(\mathcal{F}) = |E(T_{\mathcal{F}})|$, the number of edges in a spanning forest of $\Delta_1(\mathcal{F})$.*

A *face* of $P_{\mathcal{F}}$ is a subset of $P_{\mathcal{F}}$ on which a linear function is maximized. A vector $c = (c_1, \dots, c_r) \in \mathbb{R}^r$ defines a partition $C = (C_1, C_2, \dots, C_s)$ of $[r]$ into nonempty subsets, so that $c_{i_1} = c_{i_2}$ when i_1 and i_2 are in the same part of the partition, and $c_{i_1} < c_{i_2}$ whenever $i_1 \in C_{\ell_1}, i_2 \in C_{\ell_2}, \ell_1 < \ell_2$. Then the points of the face Q that maximizes $c^T x$ satisfy the equations

$$\sum_{i \in C_{\ell}} x_i = |\{F \in \mathcal{F} : F \cap C_{\ell} \neq \emptyset, F \cap C_m = \emptyset \text{ for } m > \ell\}|.$$

for $\ell = 1, 2, \dots, s$. The face that maximizes $c^T x$ is therefore the Minkowski sum of the simplices in the family

$$\mathcal{F}^C := \{F \cap C_{\ell_F} : F \in \mathcal{F}, F \cap C_{\ell_F} \neq \emptyset, F \cap C_m = \emptyset \text{ for } m > \ell_F\}$$

The dimension of the face is determined by the number of connected components of the simplicial complex $\Delta_{\mathcal{F}^C}$. If $\Delta_{\mathcal{F}^C}$ is obtained from $\Delta_{\mathcal{F}}$ by splitting one of the components of $\Delta_{\mathcal{F}}$ in two, then the corresponding face of $P_{\mathcal{F}}$ is a facet, and the coefficients of the vector c corresponding to C can be assumed to be 0 and 1. Therefore, all facets of $P_{\mathcal{F}}$ are of the form $\sum_{i \in D} x_i = t$ for some subset D of $[r]$ and integer t . When $\Delta_{\mathcal{F}^C}$ has exactly one component of size two, say $\{i, j\}$, and otherwise all isolated elements, then the corresponding face of $P_{\mathcal{F}}$ is an edge parallel to $e_i - e_j$. Vertices of $P_{\mathcal{F}}$ are points that maximize linear functions $c^T x$ in which all components of c are distinct. If $c_1 < c_2 < \dots < c_r$ then component v_i of the vertex that maximizes $c^T x$ equals the number of sets F for which i is the largest element. In particular, vertices of $P_{\mathcal{F}}$ have integer coordinates.

2 Minkowski sum of a fixed number of simplices

Suppose that \mathcal{F} consists of k subsets F_1, F_2, \dots, F_k of $[r]$. For each $i \in [r]$, define $N_{\mathcal{F}}(i) = \{j \in [k] : i \in F_j\}$. Let A be a subset of $[r]$ so that $N_{\mathcal{F}}(i_1) = N_{\mathcal{F}}(i_2)$ whenever i_1 and i_2 are in A . We would like to show how the combinatorial type of $P_{\mathcal{F}}$ can be inferred from that of $P_{\mathcal{F}'}$, where \mathcal{F}' is




Figure 1: A sum of two triangles

obtained from \mathcal{F} by replacing each appearance of A in a set F by the one-element set $m = \max(A)$. Afterward, we will restrict our attention to families in which all of the $N_{\mathcal{F}}(i)$ are distinct.

Every point $y \in P_{\mathcal{F}'}$ corresponds to the simplex $\Delta(y) := \{z \in \mathbb{R}^r : z_i = y_i, i \notin A, \sum_{i \in A} z_i = y_m, z_i \geq 0, i \in A\}$ contained in $P_{\mathcal{F}}$. Note that $\Delta(y)$ is $(|A| - 1)$ -dimensional if $y_m > 0$ and a point otherwise. Let \mathcal{F}'' be the face of \mathcal{F} where $y_m = 0$. The combinatorial type of $P_{\mathcal{F}}$ is therefore that of $\Delta_A \times P_{\mathcal{F}'}$, with (if $P_{\mathcal{F}''}$ is nonempty) the face $\Delta_A \times P_{\mathcal{F}''}$ collapsed to a copy of $P_{\mathcal{F}''}$. In the case that $|A| = 2$, $P_{\mathcal{F}}$ is a wedge over $P_{\mathcal{F}'}$ with foot $P_{\mathcal{F}''}$.

EXAMPLE Consider the family $\mathcal{F} = \{\{1, 2, 3\}, \{1, 2, 4\}\}$ of subsets of $[4]$. Then $N_{\mathcal{F}}(i) = \{1, 2\}$ for all i in $A = \{1, 2\}$. The polytope $P_{\mathcal{F}}$ is drawn in Figure 1. The polytope $P_{\mathcal{F}'}$ is the two-dimensional cube that is the top face of the drawing. $P_{\mathcal{F}''}$ is the vertex $(0, 0, 1, 1)$.

Proposition 2.1 *Every vertex of $P_{\mathcal{F}}$ is of the form $y + y_m(e_i - e_m)$, where y is a vertex of $P_{\mathcal{F}'}$ and $i \in \{1, 2, \dots, m\}$. Vertices $y + y_m(e_i - e_m)$ and $y' + y'_m(e_j - e_m)$ of $P_{\mathcal{F}}$ are adjacent in $P_{\mathcal{F}}$ if*

1. $y = y'$ and $y_m > 0$ or
2. y is adjacent to y' in $P_{\mathcal{F}'}$ and either $i = j$ or $y_m y'_m = 0$.

Note that no vertex has more than one component of A nonzero, because the vertices of the simplex Δ_A have only one nonzero component.

We consider first the case in which \mathcal{F} consists of two sets, F and F' . In the special case where each of the sets $F \setminus F'$, $F \cap F'$ and $F' \setminus F$ has exactly one element, say 1, 2 and 3 respectively, then $F = \{1, 2\}$ and $F' = \{2, 3\}$ and the Minkowski sum $P = \Delta_F + \Delta_{F'}$ is the convex hull of $(1, 1, 0)$, $(0, 2, 0)$, $(0, 1, 1)$ and $(1, 0, 1)$ in \mathbb{R}^3 , which constitutes a two-dimensional rhombus within the positive octant of the plane $x + y + z = 2$.

We now argue that the generic Minkowski sum of two simplices roughly has the structure of such a rhombus, if each of $F \setminus F'$, $F \cap F'$, and $F' \setminus F$ is nonempty.

By assigning the 1st, 2nd and 3d coordinate axis of \mathbb{R}^3 to these parts respectively, we can partition the vertices of $P = \Delta_F + \Delta_{F'}$ in the following “rhombus”-way: A vertex $e_i + e_j$ of $P_{\mathcal{F}}$ is of type $A = (1, 1, 0)$ if $i \in F \setminus F'$ and $j \in F \cap F'$, of type $B = (0, 2, 0)$ if $i = j \in F \cap F'$, of type $C = (0, 1, 1)$ if $i \in F \cap F'$ and $j \in F' \setminus F$ and of type $D = (1, 0, 1)$ if $i \in F \setminus F'$ and $j \in F' \setminus F$. Note that the rhombus formed by A, B, C and D in \mathbb{R}^3 has edges AB, BC, CD and DA . With this setup we have the following.

Lemma 2.2 *If A, B, C and D are the points in \mathbb{R}^3 as here above and $F \setminus F'$, $F \cap F'$ and $F' \setminus F$ are all nonempty, then there are no AC nor BD type edges of $P = \Delta_F + \Delta_{F'}$.*

Proof. The original rhombus does not have AC or BD edges. \square

By the above Lemma 2.2 we have the following corollary that describes the structure of a Minkowski sum of two standard simplices to be roughly that of the rhombus mentioned above.

Corollary 2.3 *If $F, F' \subseteq [r]$ then the edges, or one-dimensional faces, of $P = \Delta_F + \Delta_{F'}$ are of the following types:*

1. *Internal XX edges, where both the endvertices are of type $X \in \{A, B, C, D\}$.*
2. *XY edges, with $XY \in \{AB, BC, CD, DA\}$, where one endvertex is of type X and the other of type Y .*

Theorem 2.4 *Let $F, F' \subseteq [r]$ and let u be a vertex of the polytope $P_{\mathcal{F}}$.*

1. *If u is of type A, B or C , then $\deg(u) = |F \cup F'| - 1$.*
2. *If u is of type D , then $\deg(u) = |F| + |F'| - 2$.*

Proof. If u is of type B , say $u = 2e_i$, then u is adjacent to all $|F \cap F'| - 1$ other vertices of type B , and all type A and C vertices of the form $e_i + e_j$, where $j \in (F \setminus F') \cup (F' \setminus F)$. If u is of type A , say $u = e_i + e_j$, with $i \in F \setminus F'$ and $j \in F \cap F'$, then u is adjacent to two kinds of type A vertices: $|F \cap F'| - 1$ vertices $e_i + e_k$ with $k \in (F \cap F') \setminus \{j\}$ and $|F \setminus F'| - 1$ vertices $e_k + e_j$ with $k \in F \setminus (F' \cap \{i\})$. Also, u is adjacent to $|F' \setminus F|$ type D vertices $e_i + e_k$ with $k \in F' \setminus F$, and finally u is adjacent to the vertex $2e_j$. If u is of type D , say $u = e_i + e_j$ with $i \in F \setminus F'$ and $j \in F' \setminus F$, then u is adjacent to $|(F \setminus F') \cup (F' \setminus F)| - 2$ vertices of type D obtained by replacing either e_i or e_j by an e_k for $k \in (F \setminus F') \cup (F' \setminus F)$, and u is adjacent to $|F \cap F'|$ vertices of each type A and C , obtained by replacing e_i or e_j by an e_k for $k \in F \cap F'$. \square

Corollary 2.5 *Let $F, F' \subseteq [r]$ and $P = \Delta_F + \Delta_{F'}$.*

1. *The total number of vertices of P is $|F| \cdot |F'| - |F \cap F'|(|F \cap F'| - 1)$.*
2. *The total number of one-dimensional faces (edges) of $P = \Delta_F + \Delta_{F'}$ is given by*

$$\frac{1}{2} \left[|F \setminus F'| \cdot |F' \setminus F|(|F| + |F'| - 2) + |F \cap F'|(|F \cup F'| - 1)(|F \setminus F'| + |F' \setminus F| + 1) \right].$$

Proof. The number of vertices of degree $|F| + |F'| - 2$ in P is $|F \setminus F'| \cdot |F' \setminus F|$. By Theorem 2.4 the remaining vertices of P all have degree $|F \cup F'| - 1$. By the Hand-Shaking Theorem the total number of edges, or one-dimensional faces, is given as stated. \square

Assuming that $F \cup F' = [r]$, then the maximum value of $|F| + |F'| - 2$ (provided $F \setminus F'$ and $F' \setminus F$ are nonempty) is $2r - 4$, which occurs when $F = [r - 1]$ and $F' = [r] \setminus \{1\}$. Considering the distribution of the two possible degrees of a Minkowski sum of two simplices $P = \Delta_F + \Delta_{F'}$, we have the following.

Proposition 2.6 *Let $r \in \mathbb{N}$ be fixed. If $F, F' \subseteq [r]$ and $P = \Delta_F + \Delta_{F'}$ is of dimension $r - 1$, then the average degree $\overline{\deg}(P)$ satisfies*

$$r - 1 \leq \overline{\deg}(P) < \frac{10}{9}(r - 1).$$

Moreover, the lower bound is attained iff (i) $F \subseteq F'$, (ii) $F' \subseteq F$ or (iii) $|F \cap F'| = 1$. Also, $\overline{\deg}(P)/(r - 1)$ can become arbitrarily close to $10/9$ for large r .

Proof. We introduce the variables x, y and z by $x = |F \setminus F'|$, $y = |F' \setminus F|$ and $z = |F \cap F'|$. Here we have the boundary condition $x, y \geq 0$ and $x + y + z = r$, and since P is assumed to have dimension $r - 1$ we have $z \geq 1$ or $0 \leq x + y \leq r - 1$. By Corollary 2.5 and the Hand-Shaking Theorem we obtain that

$$\begin{aligned} \overline{\deg}(P) &= 2 \frac{|E(\Delta_1(\mathcal{F}))|}{|V(\Delta_1(\mathcal{F}))|} \\ &= \frac{|F \setminus F'| \cdot |F' \setminus F| (|F| + |F'| - 2) + |F \cap F'| (|F \cup F'| - 1) (|F \setminus F'| + |F' \setminus F| + 1)}{|F| \cdot |F'| - |F \cap F'| (|F \cap F'| - 1)} \\ &= \frac{xy(2r - 2 - x - y) + (r - 1)(r - x - y)(x + y + 1)}{(r - y)(r - x) - (r - x - y)(r - x - y - 1)}. \end{aligned}$$

As a function of x and y we note that $\overline{\deg}(P) = \overline{\deg}(x, y)$ is symmetric, has the value of $r - 1$ on the boundary of the triangle bounded by $x = 0$, $y = 0$ and $x + y = r - 1$. By Theorem 2.4 the value $\overline{\deg}(x, y)$ is strictly larger than $r - 1$ inside the triangle. If the maximum value of $\overline{\deg}(x, y)$ is $\overline{\deg}_{\max}(r)$, then $(10r - 13)/9 < \overline{\deg}_{\max}(r) < 10(r - 1)/9$, but $\overline{\deg}_{\max}(r) - (10r - 13)/9$ tends to zero when r tends to infinity. \square

REMARK: In fact, for any $\epsilon > 0$ there is an r_0 such that for any $r \geq r_0$ we have

$$r - 1 \leq \overline{\deg}(P) < \frac{10r - 13}{9} + \epsilon.$$

The f -polynomial $f_P(q)$ of a d -dimensional polytope P is $\sum_{i=0}^d f_i q^i$, where f_i is the number of i -dimensional faces of P . Postnikov [9] shows that $f_{P \times Q}(q) = f_P(q) f_Q(q)$ and gives an elegant formula for $f_{P_{\mathcal{F}}}(q)$ in the case that \mathcal{F} is a building set. If we assume that A , \mathcal{F}' and \mathcal{F}'' are as in the discussion preceding Proposition 2.1, the f -polynomial can be decomposed as follows:

Proposition 2.7 $f_{P_{\mathcal{F}}}(q) = f_{\Delta_A}(q) f_{P_{\mathcal{F}'}}(q) - f_{\Delta_A}(q) f_{P_{\mathcal{F}''}}(q) + f_{P_{\mathcal{F}''}}(q).$

In the Example, $f_{P_{\mathcal{F}}}(q) = 7 + 11q + 6q^2 + q^3 = (2 + q)(4 + 4q + q^2) - (2 + q)(1) + 1.$

If $P_{\mathcal{F}}$ is the sum of two simplices Δ_F and $\Delta_{F'}$, then one can easily check that $P_{\mathcal{F}} = \Delta_F \times \Delta_{F'}$ when $|F \cap F'|$ is 0 or 1. This allows us to describe the f -polynomials of sums of two simplices quite easily, using the proposition with $A = F \cap F'$.

Corollary 2.8 *If $\mathcal{F} = \{F, F'\}$, where $F \cap F' = \{1, 2, \dots, m\}$, then*

$$f_{P_{\mathcal{F}}}(q) = f_{\Delta_{F \cap F'}}(q) f_{\Delta_{(F \cup m) \times \Delta_{(F' \cup m)}}}(q) - f_{\Delta_{F \cap F'}}(q) f_{\Delta_F \times \Delta_{F'}}(q) + f_{\Delta_F \times \Delta_{F'}}(q)$$

In particular, the number of vertices of $P_{\mathcal{F}}$ is $|F \cap F'|(|F \setminus F'| + 1)(|F' \setminus F| + 1) - |F \cap F'| |F \setminus F'| |F' \setminus F| + |F \setminus F'| |F' \setminus F| = |F \cap F'|(|F \setminus F'| + |F' \setminus F| + 1) + |F \setminus F'| |F' \setminus F|$ which is consistent with Corollary 2.5.

We will now generalize the results that we obtained for the sum of two simplices to larger sums.

Definition 2.9 *For $k \in \mathbb{N}$ let $\mathcal{H}(k)$ be the family of k subsets of $[2^k - 1]$ so that for $i = 1, 2, \dots, 2^k - 1$, $N_{\mathcal{H}(k)}(i)$ is the i^{th} (in lexicographic order) nonempty subset of $[k]$. Then $P(k) := P_{\mathcal{H}(k)}$ is called the k^{th} master polytope.*

Definition 2.10 *Let $\mathcal{F} = (F_1, \dots, F_k)$ and let u be a point in $P_{\mathcal{F}}$. Then $h_{\mathcal{F}}(u)$ is the point v in $P(k)$ for which, for $i = 1, 2, \dots, 2^k - 1$, we set*

$$v_i = \begin{cases} \sum_{j: N_{\mathcal{F}}(j) = N_{\mathcal{H}(k)}(i)} u_j & \text{if there is a } j \text{ with } N_{\mathcal{F}}(j) = N_{\mathcal{H}(k)}(i), \\ 0 & \text{otherwise} \end{cases}$$

REMARK: Another way to look at $v = h_{\mathcal{F}}(u)$ is as follows: For $\mathcal{F} = (F_1, \dots, F_k)$ let u be a point in $P_{\mathcal{F}}$ for which $u_i u_j > 0$ implies $N_{\mathcal{F}}(j) \neq N_{\mathcal{F}}(i)$. Then let $h_{\mathcal{F}}(u)$ be the point v in $P(k)$ where $v_{\ell_i} = u_i$ where ℓ_i is the unique element in $[2^k - 1]$ with $N_{\mathcal{H}(k)}(\ell_i) = N_{\mathcal{F}}(i)$ for each $i \in [r]$.

Theorem 2.11 *For $\mathcal{F} = (F_1, \dots, F_k)$ the point $u \in P_{\mathcal{F}}$ is a vertex of $P_{\mathcal{F}}$ if, and only if, the following conditions are met.*

1. *Each instance of $u_{i_{\alpha}} u_{i_{\beta}} > 0$, $N_{\mathcal{F}}(i_{\alpha}) = N_{\mathcal{F}}(i_{\beta})$ implies that $i_{\alpha} = i_{\beta}$.*
2. *$h_{\mathcal{F}}(u)$ is a vertex of the polytope $P(k)$.*

Proof. (Theorem 2.11 Sketch) For a point $u = e_{i_1} + \dots + e_{i_k}$ of $P_{\mathcal{F}}$ we first note that if $N_{\mathcal{F}}(i_{\alpha}) = N_{\mathcal{F}}(i_{\beta})$ and $i_{\alpha} \neq i_{\beta}$, then $u = (v + w)/2$ where v and w are the points of $P_{\mathcal{F}}$ obtained from u on one hand by replacing i_{α} by i_{β} to get v and on the other hand by replacing i_{β} by i_{α} to get w . Hence, the first condition is necessary.

Assume that u satisfies the first condition and that $h_{\mathcal{F}}(u)$ is an extreme point of $P(k)$. Since there is a supporting hyperplane in $\mathbb{R}^{2^k - 1}$ containing $h_{\mathcal{F}}(u)$ there is a corresponding supporting hyperplane in \mathbb{R}^n containing u , showing that u is a vertex of $P_{\mathcal{F}}$.

Assume finally that u satisfies the first condition and that $h_{\mathcal{F}}(u)$ is not an extreme point of $P(k)$. In this case $h_{\mathcal{F}}(u)$ is a proper convex combination of extreme points of $P(k)$. Since the first condition is satisfied, there are corresponding points of $P_{\mathcal{F}}$, such that u is a proper (in fact the same!) convex combination of these. This completes the proof. \square

For $\mathcal{F} = (F_1, \dots, F_k)$ let A_1, \dots, A_h be the vertices of the polytope $P(k)$. Similar to the case when $k = 2$ we have the following.

Theorem 2.12 *If $\mathcal{F} = (F_1, \dots, F_k)$, then the edges, or one-dimensional faces, of $P_{\mathcal{F}}$ are of the following types:*

1. *Internal $A_i A_i$ type edges, where both the endvertices are of type A_i for some $i \in \{1, \dots, m\}$.*

2. $A_i A_j$ type edges, where $A_i A_j$ is an edge of the master polytope $P(k)$.

Proof. (Sketch) Similarly to the proof of Lemma 2.2 (although with a bit more elaborate indexing scheme) one can show that there is a supporting hyperplane in \mathbb{R}^n of P containing the vertex of type A_i and the vertex of type A_j if, and only if, there is a corresponding supporting hyperplane in \mathbb{R}^{2^k-1} of $P(k)$ containing the vertices A_i and A_j . \square

Theorems 2.11 and 2.12 both reduce the structure of $P_{\mathcal{F}} \subseteq \mathbb{R}^n$ to considerations of the master polytope $P(k) \subseteq \mathbb{R}^{2^k-1}$.

3 Function Representation of Integer Points of $P_{\mathcal{F}}$

As in the previous section, we assume that $\mathcal{F} = (F_1, \dots, F_k)$, an ordered collection of k subsets of $[r]$. A function $f : [k] \rightarrow [r]$ that satisfies $f(i) \in F_i$ for each i will be called a *rep-function*. For a rep-function f we define $u(f) := e_{f(1)} + \dots + e_{f(k)}$.

Claim 3.1 *For functions $f, g : [k] \rightarrow [m]$ we have*

1. $u(f) + u(g) = u(\min\{f, g\}) + u(\max\{f, g\})$.
2. If $f \neq g$, then $u(f) \neq u(\min\{f, g\})$.

In the case $u(f) = u(g)$, we obtain by Claim 3.1 that $u(f) = u(g) = (u(\min\{f, g\}) + u(\max\{f, g\}))/2$. Hence, if an integer point $u \in P_{\mathcal{F}}$ can be represented by two distinct functions f and g , then it is not a vertex of the type polytope $P(k)$. The interesting part is the converse.

Lemma 3.2 *If v is an integer point in $P_{\mathcal{F}}$ that is not a vertex of $P_{\mathcal{F}}$, and an edge of the smallest face containing v is parallel to $e_{i_1} - e_{i_2}$, then $P_{\mathcal{F}}$ contains the points $v + e_{i_1} - e_{i_2}$ and $v - e_{i_1} + e_{i_2}$.*

Proof. First note that $v_{i_1} \neq 0$ and $v_{i_2} \neq 0$, because otherwise all points on the smallest face containing v would satisfy $x_{i_1} = 0$ or $x_{i_2} = 0$, contradicting the assumption that there is an edge of this face parallel to $e_{i_1} - e_{i_2}$. If v is on a facet of $P_{\mathcal{F}}$ given by $\sum_{i \in T} x_i = t$ for some $T \subset [r]$ and integer t , then this equation is satisfied by all points in the smallest face containing v . That means that i_1 and i_2 are either both in or both outside of T . Thus $v + e_{i_1} - e_{i_2}$ and $v - e_{i_1} + e_{i_2}$ will satisfy any equations that v satisfies. Furthermore, any inequality $x_i \geq 0$ or $\sum_{i \in T} x_i \leq t$ that v satisfies strictly will also be satisfied by $v + e_{i_1} - e_{i_2}$ and $v - e_{i_1} + e_{i_2}$, because only one component is increased by 1 and one component is decreased by 1. \square

Lemma 3.3 *If f and g are rep-functions and $u(g) = u(f) + te_{i_1} - te_{i_2}$ for $i \neq j$ in $[r]$, then there exist rep-functions f_1, f_2, \dots, f_{t-1} so that $u(f) + le_{i_1} - le_{i_2} = u(f_l)$ for $l = 1, 2, \dots, t-1$.*

Proof. Define $G_{\mathcal{F}}$ to be the bipartite graph with vertex set $\{w_j : j \in [k]\} \cup \{v_t : t \in [r]\}$ and edges $\{(w_j, v_i)\}$ for all (i, j) with $i \in F_j$. For any rep-function h , let M_h be the set of edges (w_j, v_i) for which $h(j) = i$. For every $i \in [r] \setminus \{i_1, i_2\}$, the number of edges of M_g meeting v_i equals the number of edges of M_f meeting v_i . For every $j \in [k]$, w_j is met by exactly one edge from each of M_f and M_g . On the other hand, v_{i_1} is adjacent to t more edges of M_g than M_f , and v_{i_2} is adjacent to t more edges of M_f than M_g . There therefore exists a path P from v_{i_2} to v_{i_1} that alternates between edges of M_f and M_g . Let M^1 be the set of edges obtained from M_f by replacing the edges of M_f in the path by the edges of M_g in the path. Then, for $j = 1, 2, \dots, k$, define $f_1(j) = i$, where (w_j, v_i) is an edge of M^1 . Then $u(f_1) = u(f) + e_{i_1} + e_{i_2}$. We can continue this way to get $u(f_2), \dots, u(f_{t-1})$. \square

Proposition 3.4 *Every integer point v in $P_{\mathcal{F}}$ is $u(f)$ for some rep-function f .*

Proof. The proof is by induction on the dimension of the smallest face containing v . From the first section, we know that the statement is true if v is a vertex. Suppose v is not a vertex. Suppose that there is an edge of the smallest face containing v that is parallel to $e_{i_1} - e_{i_2}$. Then lemma 3.2 allows us to build a segment parallel to $e_{i_1} - e_{i_2}$, containing v in its interior, and with endpoints on faces of $P_{\mathcal{F}}$ that are of lower dimension than the one containing v . By induction, the endpoints of the interval are $u(f)$ and $u(g)$ for some rep-functions f and g . Lemma 3.3 then gives us a rep-function for v . \square

Theorem 3.5 *An integer point v in $P_{\mathcal{F}}$ is a vertex of $P_{\mathcal{F}}$ if and only if there is a unique rep-function f so that $u(f) = v$.*

Proof. Let v be an integer point in $P_{\mathcal{F}}$ that is not a vertex of $P_{\mathcal{F}}$. By Lemma 3.2 there are i_1 and i_2 in $[r]$ so that $P_{\mathcal{F}}$ contains the points $v - e_{i_1} + e_{i_2}$ and $v - e_{i_2} + e_{i_1}$. Let f and g be the rep-functions guaranteed by Proposition 3.4 for $v - e_{i_1} + e_{i_2}$ and $v - e_{i_2} + e_{i_1}$, respectively. Let $G_{\mathcal{F}}, M_f$ and M_g be as in the proof of Lemma 3.3. Then There are two edges of M_f adjacent to v_{i_2} that are not in M_g . Therefore we can use these edges as initial edges in two different paths from v_{i_2} to v_{i_1} that alternate between edges of M_f and M_g . Swapping edges of M_f for edges of M_g along each of these alternating paths leads to two different rep-functions for v . \square

The number of rep-functions for a given \mathcal{F} is easy to count, it is $\prod_{F \in \mathcal{F}} |F|$. By listing the rep-functions and the corresponding integer points $u(f)$, and striking out the $u(f)$ that appear more than once, one can list the vertices of $P_{\mathcal{F}}$. This was done by Bernd Sturmfels [1] for the polytopes $P(k)$, $k = 3, 4, 5$. He found that $P(3)$ had 41 vertices, $P(4)$ had 1015 vertices, and $P(5)$ had 59072 vertices.

4 Max-degree as function of parameters alone

In this section we determine the function $d : \mathbb{N} \rightarrow \mathbb{N}$ given by

$$d(r) = \max_{\mathcal{F}} \{ \deg_{\max}(P_{\mathcal{F}}) \},$$

where the maximum is taken over all multi-subsets (F_1, \dots, F_k) of $\mathbb{P}([r])$, where $k \in \mathbb{N}$ can be any integer but r is fixed. Moreover, for each fixed $k \in \mathbb{N}$ we determined the function $d_k : \mathbb{N} \rightarrow \mathbb{N}$ defined by

$$d_k(r) = \max_{|\mathcal{F}| \leq k} \{ \deg_{\max}(P_{\mathcal{F}}) \},$$

where the maximum is here taken over all multi-subsets (F_1, \dots, F_k) of $\mathbb{P}([r])$ where both k and r are fixed. Clearly $d(r) = \max_{k \in \mathbb{N}} \{d_k(r)\}$.

We start with the following lower bound for $d_k(r)$ and $d(r)$.

Lemma 4.1 *For $k, r \in \mathbb{N}$ we have $d_k(r) \geq k(r - k)$, and therefore $d(r) \geq \lfloor r^2/4 \rfloor$.*

Proof. Let $k \in [r]$ and let for each $i \in [k]$ let $F_i = \{i, k + 1, k + 2, \dots, r\}$. Then the vertex $v = e_1 + e_2 + \dots + e_k$ is adjacent to each of the vertices $v + (e_{i_1} - e_{i_2})$, for $1 \leq i_2 \leq i_1$ and $k + 1 \leq i_1 \leq r$. Therefore $d_k(r) \geq k(r - k)$, so we have in particular that $d(r) \geq \lfloor r/2 \rfloor \lceil r/2 \rceil = \lfloor r^2/4 \rfloor$. \square

Another polytope that has vertices of degree $\lfloor r^2/4 \rfloor$ is the graphical zonotope for the complete bipartite graph with $\lfloor r/2 \rfloor$ vertices on one side of the bipartition and $\lceil r/2 \rceil$ vertices on the other side. West [11] proved that the graphical zonotope for the complete bipartite graph has vertices of degree ℓ for all $r-1 \leq \ell \leq \lfloor r^2/4 \rfloor$. On the other hand, every vertex of the polytope of lemma 4.1 other than v has degree $r-1$.

For a fixed vertex u , each edge of P incident to u can be identified with a multiple of a difference $e_i - e_j$ of some pair of unit vectors, where $i, j \in [r]$ are distinct. Since the collection $\{\alpha(e_i - e_j) : \alpha \in \mathbb{N}\}$ is a set of parallel vectors, at most one multiple of $e_i - e_j$ can possibly correspond to an edge incident to u . From this alone we see that the maximum number of edges incident to u is at most $\binom{r}{2}$. However, more can be said:

For a vertex u of P , let $\vec{G}(u)$ be the directed graph with the vertexset $V(\vec{G}(u)) = [r]$ where a directed edge (i, j) is present iff $u + \alpha(e_i - e_j)$ is a neighbor of u in P for some $\alpha \in \mathbb{N}$.

Proposition 4.2 *For $r \in \mathbb{N}$ and $\mathcal{F} = (F_1, \dots, F_k) \subseteq \mathbb{P}([r])$, the digraph $\vec{G}(u)$ is acyclic and its underlying graph $G(u)$ is simple and triangle-free.*

Proof. Assume there is a cycle (i_1, i_2, \dots, i_h) in $\vec{G}(u)$. Then u, v_1, \dots, v_h are all vertices of P , where $v_\ell = u + \alpha_\ell(e_{i_\ell} - e_{i_{\ell+1}})$ (here we compute cyclically, so $e_{i_{h+1}} = e_{i_1}$). This is however impossible since

$$\sum_{\ell=1}^h \frac{1}{\alpha_\ell} (v_\ell - u) = 0,$$

which means that there is no hyperplane containing u alone and having all the v_ℓ 's strictly on one side of it. In particular for $h = 2$, there are no directed 2-cycles and hence the underlying graph $G(u)$ is simple. Also for $h = 3$, there are no directed triangles in $\vec{G}(u)$ either.

Assume now that $G(u)$ has a triangle, which then does not correspond to a directed triangle in $\vec{G}(u)$, say $v = u + \alpha(e_i - e_j)$, $v' = u + \beta(e_j - e_l)$ and $v'' = u + \gamma(e_i - e_l)$. In this case we have

$$v'' - u = \frac{\gamma}{\alpha} (v - u) + \frac{\gamma}{\beta} (v' - u),$$

which means that the vector $v'' - u$ is in the cone spanned by $v - u$ and $v' - u$. This contradicts the fact that uv'' is an edge of P . Hence, the underlying graph $G(u)$ of $\vec{G}(u)$ has no triangles. \square

Theorem 4.3 *For $r \in \mathbb{N}$ we have $d(r) \leq \lfloor r^2/4 \rfloor$.*

Proof. The maximum degree of a vertex u of P is by Proposition 4.2 the maximum number of edges the simple triangle free graph $G(u)$ can have. By a theorem by Mantel [7] (as a special case of Turán's Theorem [10]), the maximum number of edges of a simple triangle-free graph on r vertices is $\lfloor r^2/4 \rfloor$, hence the theorem. \square

By Lemma 4.1 and Theorem 4.3 we have the following corollary.

Corollary 4.4 *For $r \in \mathbb{N}$ we have $d(r) = \lfloor r^2/4 \rfloor$.*

We now turn our attention to the computation of $d_k(r)$. Note that the Minkowski sum $P_{\mathcal{F}}$ provided in the proof of Lemma 4.1 that attains the overall maximum degree $d(r)$ has $k = |\mathcal{F}| = \lfloor r/2 \rfloor$. Therefore when computing $d_k(r)$ we can assume $1 \leq k \leq r/2$.

First we need a variation of the theorem by Mantel [7]: Let G be a simple graph on n vertices and let $1 \leq k \leq n/2$.

Call G a k -triangle-free graph, or a k -tr for short, if G is triangle free and G has a vertex cover of cardinality at most k .

Theorem 4.5 *Let $n \in \mathbb{N}$ and $1 \leq k \leq n/2$. If $e_k(n)$ is the maximum number of edges of a k -tr graph G , then $e_k(n) = k(n - k)$. Moreover, if G is a k -tr graph on n vertices with $e_k(n)$ edges, then G is a complete bipartite with parts of cardinalities k and $n - k$.*

Proof. For $n \in \{1, 2\}$ the theorem is trivial. We proceed by induction and assume we have a k -tr graph on $n > 2$ vertices with the maximum number $e_k(n)$ of edges. Let $uv \in E(G)$ be an edge and since either u or v is in the vertex cover U of size k , we assume it to be u . Since G is triangle-free the set of neighbors $N(u)$ and $N(v)$ are disjoint. Let $G' = G - \{u, v\}$ be the simple graph obtained from G by removing the vertices u and v from G . By the disjointness of $N(u)$ and $N(v)$ we have $|E(G)| = |E(G')| + d(u) + d(v) - 1$.

Assume first that $v \in U$. In this case G' is a $(k - 2)$ -tr graph on $n - 2$ vertices and hence by induction hypothesis we have $|E(G)| = |E(G')| + d(u) + d(v) - 1 \leq (k - 2)[(n - 2) - (k - 2)] + n - 1 < k(n - k)$.

Now assume that $v \notin U$. In this case G' is a $(k - 1)$ -tr graph on $n - 2$ vertices and hence by induction hypothesis we have $|E(G)| = |E(G')| + d(u) + d(v) - 1 \leq (k - 1)[(n - 2) - (k - 1)] + n - 1 = k(n - k)$. Also by inducting hypothesis, $|E(G)| = k(n - k)$ can hold iff G' is a complete bipartite graph with parts of cardinalities $k - 1$ and $n - k - 1$, and $d(u) + d(v) = n$ (i.e. $N(u) \cup N(v) = V(G)$). This means that $|E(G)| = k(n - k)$ can hold iff $N(v) = U$ and $N(v) = V(G) \setminus U$, that is, G is a complete bipartite graph with parts of sizes k and $n - k$. This completes the proof. \square

From Theorem 4.5 we obtain the following corollary.

Corollary 4.6 *For $r \in \mathbb{N}$ and $k \in \{1, \dots, \lfloor r/2 \rfloor\}$, we have $d_k(r) = k(n - k)$.*

Proof. Consider a point $u = e_{i_1} + \dots + e_{i_k}$ of $P_{\mathcal{F}}$ (note that some indices might coincide). As noted before, a neighbor v of u in P must have the form $v = u + \alpha(e_i - e_j)$ for some $\alpha \in \mathbb{N}$, and $i \in [r]$ and $j \in \{i_1, \dots, i_k\}$. Since each directed edge $(i, j) \in V(\vec{G}(u))$ has its head in $\{i_1, \dots, i_k\}$, of cardinality at most k , the underlying graph $G(u)$ has a vertex cover of size at most k . Therefore $G(u)$ is a k -tr graph and hence by Theorem 4.5 at most $k(r - k)$ edges.

In the proof of Lemma 4.1 an example of $P_{\mathcal{F}}$ with $|\mathcal{F}| \leq k$ and a vertex of degree $k(n - k)$ was given. This completes the argument. \square

5 Minkowski sum of three simplices

In this section we will investigate the polytope $P(3)$. Let $\mathcal{H} := \mathcal{H}(3) = (\{1, 2, 4, 5\}, \{1, 2, 3, 6\}, \{1, 3, 4, 7\})$. Henceforth we will drop the (3). Then $N_{\mathcal{H}}(1) = \{1, 2, 3\}$, $N_{\mathcal{H}}(2) = \{1, 2\}$, $N_{\mathcal{H}}(3) = \{2, 3\}$, $N_{\mathcal{H}}(4) = \{1, 3\}$, $N_{\mathcal{H}}(5) = \{1\}$, $N_{\mathcal{H}}(6) = \{2\}$, $N_{\mathcal{H}}(7) = \{3\}$, so all of the nonempty subsets of $[3]$ are represented. The case of $k = |\mathcal{F}| = 3$ is the first interesting case for the mere reason that the polytope $P(3)$ does not have $2^{k(k-1)} = 64$ vertices, as was the case for $k = 2$, where the rhombus $P(2)$ had precisely $2^{k(k-1)} = 4$ vertices.

EXAMPLE: the point $A = (0, 1, 1, 1, 0, 0, 0)$ in $P(3)$ is not a vertex, because $A = (B + C + D)/3$, where $B = (0, 2, 1, 0, 0, 0, 0)$, $C = (0, 0, 2, 1, 0, 0, 0)$ and $D = (0, 1, 0, 2, 0, 0, 0)$ and all the points B, C and D are points in the polytope $P(3)$.

Lemma 5.1 *The polytope $P(3)$ has 41 vertices in \mathbb{R}^7 given by the column vectors (without the last entry) in the following 7×10 , 7×21 and 7×10 matrices. The last entry in each column is the degree of the vertex.*

3	1	1	0	0	1	0	0	0	0
0	0	2	2	1	0	1	2	0	0
0	2	0	1	2	0	0	0	2	1
0	0	0	0	0	2	2	1	1	2
0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0
<hr/>									
6	6	6	6	6	6	6	6	6	6

2	1	1	0	0	0	0	2	1	1	0	0	0	0	2	1	1	0	0	0	0
0	0	1	1	1	0	0	0	0	1	1	1	0	0	0	0	0	1	1	0	2
0	0	0	1	0	1	2	0	1	0	1	0	1	0	0	0	1	1	0	1	0
0	1	0	0	1	1	0	0	0	0	0	1	1	2	0	1	0	0	1	1	0
1	1	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	1	1	1	1	1	1	1	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	1	1	1
<hr/>																				
6	6	6	6	8	6	8	6	6	6	8	6	6	8	6	6	6	6	6	8	8

1	0	0	1	0	0	1	0	0	0
0	0	0	0	1	0	0	1	0	0
0	1	0	0	0	0	0	0	1	0
0	0	1	0	0	1	0	0	0	0
1	1	1	0	0	0	1	1	1	1
1	1	1	1	1	1	0	0	0	1
0	0	0	1	1	1	1	1	1	1
<hr/>									
7	8	8	7	8	8	7	8	8	9

These computations were verified using the computer program POLYMAKE [5]. Using POLYMAKE, we determined that the polytope $P(4)$ had vertices of all degrees in the set $\{14, 15, \dots, 28\}$ except for $\{16, 23, 26, 27\}$.

Acknowledgments

The authors would like to thank James F. Lawrence for helpful discussions regarding the theory of polytopes in general. Last but not least, sincere thanks to Bernd Sturmfels for introducing this problem to us and for his keen encouragement.

References

- [1] Bernd Sturmfels, Personal communication (2005).
- [2] Aldo Conca, Linear Spaces, Transversal Polymatroids and ASL Domains, [arXiv:math.AC/0504111](https://arxiv.org/abs/math.AC/0504111) v1 6 Apr 2005.

- [3] Eva Maria Feichtner and Bernd Sturmfels, Matroid Polytopes, Nested Sets and Bergman Fans, Preprint, [arXiv:math.CO/041126](#) v1 11 Nov 2004.
- [4] D. Fisher, K. Fraughnaugh, L. Langley, D. West, The Number of Dependent Arcs in an Acyclic Orientation, *Journal of Combinatorial Theory (B)* **71**, 73 – 78, (1997).
- [5] E. Gawrilow and M. Joswig, Polymake: a framework for analyzing convex polytopes, in *Polytopes - Combinatorics and Computation*, eds. G. Kalai and G. M. Ziegler, Birkhäuser, 2000, pp. 43–74.
- [6] J. Herzog and T. Hibi, Discrete Polymatroids, *Journal of Algebraic Combinatorics*, **16**, 239 – 268, (2002).
- [7] W. Mantel, Problem 28, soln. by H. Gouwentak, W. Mantel, J. Teixeira de Matters, F. Schuh and W. A. Wythoff, *Wiskundige Opgaven*, **10**, 60 – 61, (1907).
- [8] J. Morton, L. Pachter, A. Shiu, B. Sturmfels, O. Wienand Geometry of Rank Tests, Preprint, [arXiv:math.ST/0605173](#) v1 7 May 2006.
- [9] A. Postnikov, Permutohedra, Associahedra, and Beyond [arXiv:math.CO/0507163](#) v1 7 Jul 2005.
- [10] P. Turán, Eine Extremalaufgabe aus der Graphentheorie, *Mat. Fiz Labook*, **48**, 436 – 452, (1941).
- [11] D. West, Acyclic orientations of complete bipartite graphs *Discrete Mathematics*, **138** 393 – 396, (1995).

February 2, 2008